Physics at the Energy Frontier

Aaron Pierce
University of Michigan
Fermilab Users Meeting
June 5, 2008



New Physics Motivations?

- Electroweak Symmetry Breaking
 - Hierarchy Problem
- Dark Matter
- Matter--Anti-Matter Asymmetry
 - EWBG--in MSSM very well defined region
- Flavor
- GUTs
- Dark Energy???



Find the Higgs...

- The first step to understanding electroweak symmetry breaking is to find the Higgs Boson.
- An experimental triumph, but will be unlikely to drastically affect how we think about the weak scale.





Beyond the Standard Model at the Weak Scale

- Hierarchy Problem
- Dark Matter
- In specific scenarios, there are other motivations, but these are generic.



Natural Theories?

$$\mathcal{L} \supset -m_h^2 |h|^2 + \frac{\lambda}{4} |h|^4$$

• Hierarchy Problem:

$$\Delta m_h^2 = \frac{3y_{top}^2}{8\pi^2} \Lambda^2$$

$$m_h^2 = (m_h^2)_{bare} + \text{correction}$$

Natural Theories?

Hierarchy Problem:

Motivates most BSM Physics:

- Supersymmetry
 - Technicolor
 - Little HiggsLED
- •Randall-Sundrum



Naturalness?

- Two tales:
 - Electron Classical Radius
 - Cosmological Constant



E&M Correction to Electron

$$\Delta E_{Coulomb} = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r_e}$$

$$m_e c^2 = (m_e c^2)_{bare} + \Delta E_{Coulomb}$$

0.511 = -999999.489 + 100000.000 MeV

H. Murayama hep-ph/0002232



Positron to the rescue.

To avoid fine tuning, need something new at distances of order 10-13cm

$$\Delta E = \Delta E_{Coulomb} + \Delta E_{pair}$$

$$= \frac{3\alpha}{4\pi} m_e c^2 \log \frac{\hbar}{m_e c r_e}$$

$$m_e c^2 = (m_e c^2)_{bare} (1 + \frac{3\alpha}{4\pi} \log \frac{\hbar}{m_e c r_e})$$



Pion Mass Splitting

$$m_{\pi^{\pm}}^2 - m_{\pi^0}^2 = \frac{3\alpha}{4\pi} \Lambda^2$$

Here, the cutoff is supplied by the mass of the rho meson.



Cosmological Constant

- Λ≈(meV)⁴
- This represents a fine-tuning of 120 orders of magnitude.
- Where is the new physics at the meV scale?
- Do we really understand fine-tuning?




Will we see positron/pion story again?

 If so, then we are likely to have a host of new particles responsible for softening the

divergences again.

Supersymmetry. (stops)

Little Higgs.



G. Azuelos et al. Published in Eur.Phys.J.C39S2:13-24,2005



Naturalness?


If not, what can we expect at the TeV scale? We should think about things different from Little Higgs/Technicolor/SUSY.

But What?



Dark Matter

- Second Motivation For New Physics
 Beyond the Standard Model at the Weak
 Scale:
 - Thermal Relic Abundance of Dark Matter





Dark Matter

$$\Omega_{DM} = \frac{(4\pi)^2}{3} \sqrt{\frac{\pi}{45}} \frac{x_f g_S(\gamma)}{g_*^{1/2}} \frac{T_{\gamma}^3}{H_0^2 M_P^3 \sigma} = 0.22 \frac{\text{pb}}{\sigma}$$

- So the exquisite mesurment of the Dark Matter Density (due to WMAP and others) points to a weak scale cross section.
- We already think the weak scale is interesting from a particle physics point of view. A cruel coincidence?



Does Dark Matter Guarantee Discoveries at Tevatron/LHC?


- No.
 - Simplest possibility: Dark Matter is nonthermal. Lee-Weinberg discussions is a coincidence.
 - Axion.
- What if Lee-Weinberg right?



 A simple possibility that won't cause the champagne to flow in Geneva:

 An n-plet charged under SU(2) with an X⇔-X symmetry.

Gauge Interactions Allow for Annihilation



Strumia, Cirelli, Fornengo, Tamburini



Morale:

- Dark Matter by itself won't ensure exiciting phenomenology.
- SUSY-- good dark matter candidate.
- If Dark Matter is part of a larger structure, that has TeV colored particles, then we are in business. Structure could be SUSY/UED/ Little Higgs models designed to solve hierarchy problem. What else?



SUSY Scorecard

Successes

Gauge Coupling Unification

Dark Matter

Failures

Superpartners?

Higgs Mass

FCNC (b to s gamma)

EDMS?

proton decay (dim 5)


Gravitinos?

Higgs Fine Tune


CC



Trouble with Scalars



e.g. Gabbiana, et a hep-ph/9604387



Murayama and AP hep-ph/0108104



Gauge Coupling Unification

- Scalars (and SM fermions) come in complete GUT multiplets don't contribute to relative running
- Higgsinos, Gauginos, Gauge Bosons, Higgs Bosons are responsible for unification.



Split Supersymmetry

- No attempt is made to solve the gauge hierarchy problem.
- Prejudice: Supersymmetry has something to do with a fundamental theory.
- New Structure (SUSY) + Dark
 Matter⇒Weak Scale Gauginos

Dimopoulos, Arkani-Hamed; Giudice, Romanino




Mass Spectrum






Split SUSY

- Not only electroweakinos, but Gluinos!
- Due to heavy squark mass, these can have a long lifetime, but likely less than a ~second due to BBN constraints. (Arvanitaki, Davis, Graham, AP, Wacker)





$$\frac{dE}{dx} = -\frac{4\pi\alpha^2\rho}{Am_p m_e} \frac{Zz^2}{v^2} \left(\frac{1}{2} \ln \frac{2m_e v^2}{I(1-v^2)} - v^2 \right)$$



Stopping Gluinos




Stopping Gluinos

- Stopped Gluinos show up in the detector as a jet--- but not from the origin.
- Gluinos are pair produced. About 20% of the time that one gluino stops, they both do.
- Separation between "bangs" give us the gluino lifetime.

Also staus as NLSP: Feng/ Smith, Ellis/R, /Raklev/Oye, ...



A Tevatron Result




hep-ph/07050306 (DZero)



Is the weak scale finetuned?

Most have a top-partner.

$$\Delta m_h^2 = \frac{3y_{top}^2}{8\pi^2} \Lambda^2$$



• Examples exist (e.g., Thaler and AP) where you see a top partner, (gg to TT to tops + missing) but the Higgs is still fine-tuned.



Conclusions

- Tevatron data rolling in-- surprises?
- LHC should deliver us the particle responsible for electroweak symmetry breaking.
 - Naturalness-- implications for understanding CC?
 - Dark Matter.

